

Mean-field phase diagram of interacting e_g electrons

Marcin Raczkowski,^{a,b,*} Andrzej M. Oleś^b Raymond Frésard^a

^aLaboratoire CRISMAT, UMR CNRS-ENSICAEN(ISMRA) 6508, 6 Bld. du Maréchal Juin, F-14050 Caen, France

^bMarian Smoluchowski Institute of Physics, Jagellonian University, Reymonta 4, PL-30059 Kraków, Poland

Abstract

We investigate the magnetic phase diagram of the two-dimensional model for e_g electrons which describes layered nickelates. One finds a generic tendency towards magnetic order accompanied by orbital polarization. For two equivalent orbitals with diagonal hopping such orbitally polarized phases are induced by finite crystal field.

Key words: layered nickelates, degenerate Hubbard model, orbital order

PACS: 75.10.Lp, 71.10.Fd, 71.30.+h

Correlated e_g electrons lead to several interesting phenomena in cuprates and manganites. At large intraorbital Coulomb interaction U and finite Hund's exchange J_H , the phase diagram at the filling $n = 1$ electron per site shows an interesting competition between ferromagnetic (FM) and antiferromagnetic (AF) instabilities [1]. Recent studies have further emphasized the role of J_H at orbital degeneracy [2]; here we address this problem starting from weak coupling.

We consider a two-dimensional (2D) model,

$$H = H_0 + H_{int}, \quad (1)$$

for e_g electrons, with the one-particle term:

$$H_0 = \sum_{\langle ij \rangle} \sum_{\alpha\beta\sigma} t_{ij}^{\alpha\beta} c_{i\alpha\sigma}^\dagger c_{j\beta\sigma} + \frac{1}{2} E_z \sum_i (n_{ix} - n_{iz}). \quad (2)$$

For e_g orbitals: $|x\rangle \sim |x^2 - y^2\rangle$ and $|z\rangle \sim |3z^2 - r^2\rangle$, one has the hopping matrix $t_{ij}^{\alpha\beta} = -\frac{t}{4} \begin{pmatrix} 3 & \pm\sqrt{3} \\ \pm\sqrt{3} & 1 \end{pmatrix}$, where t stands for a ($dd\sigma$) hopping matrix element, and the off-diagonal hopping along a and b axis depends on the phases of the $|x\rangle$ orbital. The crystal field splitting $\propto$

E_z , *a priori* finite in a 2D model, lifts the degeneracy of e_g orbitals. We compare this case with the frequently studied *diagonal-hopping model* [2], $t_{ij}^{\alpha\beta} = -\frac{t}{2} \delta_{\alpha\beta}$, i.e., with the degenerate Hubbard model.

Coulomb interactions H_{int} are described by on-site intraorbital Coulomb U and exchange J_H elements [3]; here we write the leading terms using the operators: $n_i = \sum_{\alpha\sigma} n_{i\alpha\sigma}$, $m_i = \sum_{\alpha\sigma} \lambda_\sigma n_{i\alpha\sigma}$, $o_i = \sum_{\alpha\sigma} \lambda_\alpha n_{i\alpha\sigma}$, and $p_i = \sum_{\alpha\sigma} \lambda_\alpha \lambda_\sigma n_{i\alpha\sigma}$, with $\lambda_\alpha = \pm 1$ for $\alpha = x(z)$ orbital, and $\lambda_\sigma = \pm 1$ for $\sigma = \uparrow(\downarrow)$ spin:

$$H_{int} = \frac{1}{8} \sum_i [(3U - 5J_H)n_i^2 - (U + J_H)m_i^2 - (U - 5J_H)o_i^2 - (U - J_H)p_i^2]. \quad (3)$$

For the electron density $n < 2$ (doping $x = 2 - n$), the model (1) describes layered $\text{La}_{2-x}\text{Sr}_x\text{NiO}_4$ nickelates. The coefficient $(U + J_H)$ of the m_i^2 term is the Stoner parameter.

We investigate the stability of various phases with either uniform or staggered magnetic order by tackling the above Hamiltonian in mean-field approximation on a 128×128 cluster, using periodic boundary conditions at low temperature $T = 0.01t$, for two representative values of Hund's exchange: $J_H = 0.15U$ and $J_H = 0.25U$. One finds that the paramagnetic (PM) phase can be unstable towards: (i) orbitally polarized paramagnetic ($\text{PM}\alpha$) phase (with the enhanced elec-

* Corresponding author (Marcin Raczkowski).

Email address: m.raczkowski@if.uj.edu.pl

¹ M.R. was supported by a Marie Curie fellowship of the European Community program under number HPMT2000-141. This work was supported by the Polish State Committee of Scientific Research (KBN) under Project No. 1 P03B 068 26.




Fig. 1. Phase diagrams of the e_g orbital model (1) as functions of the Stoner parameter $U + J_H$ and hole doping $x = 2 - n$, with $E_z = 0$ (a,b) and $E_z = 0.1U$ (c,d). Left (right) panels show results for $J_H = 0.15U$ ($J_H = 0.25U$). Dashed (dotted) line indicates the AFx-PMz (PMx-PMz) phase transition.

tron density in α orbitals); (ii) FM or orbitally polarized FM (FM α) phases; (iii) AF or orbitally polarized AF (AF α) ones. The phases with orbital polarization are generic for the 2D model of e_g electrons (Fig. 1) – they are characterized by finite values of the $\langle o \rangle$ and possibly $\langle p \rangle$ order parameters. For instance, both types of AF phases, AF x with $\langle o \rangle > 0$ (the system favors $|x\rangle$ over $|z\rangle$ occupancy), and AF z , compete with each other and are stable in different regions.

In the intermediate correlated regime $U + J_H \simeq 2t$ the AF x phase is stabilized due to the larger hopping element between $|x\rangle$ orbitals, whereas in the strongly correlated regime $U + J_H \geq 5t$, the system favors instead $|z\rangle$ orbitals which allows to better optimize the magnetic energy owing to a negative $\langle p \rangle$. However, upon increasing x the double occupancy energy becomes less important compared to a possible kinetic energy gain when $|x\rangle$ orbitals are closer to half filled. This leads to a first order phase transition at $x \gtrsim 0.6$, with all the order parameters changing discontinuously.

In a similar way one can understand the origin of PM x state (optimizing the kinetic energy) and the stability of FM z phase in the strongly correlated regime at finite doping (optimizing the magnetic energy). The latter phase is strongly favored by a large value of $J_H = 0.25U$, so that the usual AF instability at $x = 1$ is fully suppressed. Moreover, at $J_H > 0.2U$ the attractive interaction in the o channel changes into a repulsive one. As a consequence, the energy of the AF x phase cannot be much reduced by a large positive $\langle o \rangle$, and electrons redistribute almost equally over the two orbitals. Therefore, double occupancies might be avoided *even*




Fig. 2. Phase diagrams as in Fig. 1 for the degenerate Hubbard model, for: (a) $E_z = 0$ and $J_H = 0.25U$; (b) $E_z = 0.1U$, with $J_H = 0.15U$ (solid line) and $J_H = 0.25U$ (dashed line).

without creating a finite magnetic moment which makes the PM x region broader for $J_H = 0.25U$ than for $J_H = 0.15U$. We note that the realistic value of $E_z = 0.1U$, as reported for La_2NiO_4 [4], has only little influence on the phase diagram, except for strong reduction of the region of stability of the AF x phase. Remarkably, the FM z -PM x line [Fig. 1(d)] is nearly unaltered by finite E_z , showing that the FM instability almost decouples from the orbital polarization (when both $\langle o \rangle$ and $\langle p \rangle$ order parameters are suppressed).

Finally, we investigate the magnetic instabilities of the degenerate Hubbard model in the subspace with $\langle o \rangle = \langle p \rangle = 0$ [Fig. 2(a)]. In qualitative agreement with Ref. [3], the instabilities of the PM phase towards an AF/FM one for small/large doping depend only on the Stoner parameter $U + J_H$; they were also found in the strongly correlated regime [5]. The phase diagram at finite E_z [Fig. 2(b)] depends on J_H in a non-trivial way, and we recover the AF z phase in the vicinity of $x = 1$, provided J_H is small enough. Namely, when one orbital is sufficiently favored over the other one, the $S = \frac{1}{2}$ AF z phase is obtained, since some processes of the FM superexchange are blocked.

Summarizing, we have found that both the realistic e_g hopping $t_{ij}^{\alpha\beta}$ and finite crystal field splitting E_z act to stabilize *orbitally polarized phases*. The phase diagram of the e_g electron model is richer than that of the degenerate Hubbard model [3], and favors AF phases. For instance, at a fixed value of the Stoner parameter $U + J_H$, the crossover from the AF to FM phase occurs for e_g electrons at higher doping x .

References

- [1] L.F. Feiner, A.M. Oleś, and J. Zaanen, Phys. Rev. Lett. 78 (1997) 2799.
- [2] A. Klejnberg and J. Spalek, Phys. Rev. B 61 (2000) 15542; S. Florens *et al.*, *ibid.* 66 (2002) 205102; Y. Ōno, M. Potthoff, and R. Bulla, *ibid.* 67 (2003) 035119; A. Koga *et al.*, Phys. Rev. Lett. 92 (2004) 216402.

- [3] A.M. Oleś, Phys. Rev. B 28 (1983) 327.
- [4] P. Kuiper *et al.*, Phys. Rev. B 44 (1991) 4570; P. Kuiper *et al.*, *ibid.* 57 (1998) 1552.
- [5] J. Bünemann, W. Weber, and F. Gebhard, Phys. Rev. B 57 (1998) 6896.